

***Morganella austromontana* sp. nov. from the South Brazilian Plateau**

CAMILA R. ALVES ^{1*}, VAGNER G. CORTEZ ² & ROSA M. B. DA SILVEIRA¹

¹ Programa de Pós-graduação em Botânica, Universidade Federal do Rio Grande do Sul,
Av. Bento Gonçalves 9500, Agronomia CEP 91501-970, Porto Alegre, RS, Brazil

² Programa de Pós-graduação em Botânica, Departamento de Biodiversidade,
Rua Pioneiro 2153, Jardim Dallas CEP 85950-000, Palotina, PR, Brazil

* CORRESPONDENCE TO: cra.micologia@gmail.com

ABSTRACT—A new *Morganella* species was collected during field exploration of the mycobiota of the South Brazilian Plateau. It is proposed here as *Morganella austromontana*, distinguished by its ovoid basidiospores and its soil habitat. A complete description and illustrations of its morphological features are accompanied by a taxonomic discussion.

KEY WORDS—Agaricaceae, Agaricomycetidae, *Araucaria angustifolia* forest, puffballs, taxonomy

Introduction

Morganella Zeller is diagnosed by small gasteroid basidiomata (puffballs) that are globose, subglobose or depressed enclosed within a double peridium comprising a granular, verrucose, velutinous or spiny exoperidium and an endoperidium with an apical pore; basidiospores are globose or subglobose (rarely ovoid), a eucapillitium is absent, and a paracapillitium is present (Zeller 1948, Kreisel & Dring 1967). About 20 species have been included in the genus. From Brazil, 12 species have been reported and/or described in the last decade (Cortez et al. 2007; Trierveiler-Pereira & Baseia 2009; Barbosa et al. 2011; Alfredo et al. 2012, 2014; Alfredo & Baseia 2014; Alves & Cortez 2013). These puffballs occur in diverse Brazilian biome habitats, including arid environments (Barbosa et al. 2011) and seasonal ombrophilous forests (Alfredo et al. 2012, Alves & Cortez 2013) where they grow on rotting wood among litter and mosses.

During a survey of the gasteroid mycobiota in South Brazilian forests in the Atlantic Forest biome, a new *Morganella* species was found growing in the mixed ombrophilous forest typical of south Brazil. Here we describe this species as *Morganella austromontana*.

Materials & methods

Specimens were gathered during field trips in the natural reserves of the Rio das Furnas in the municipality of Alfredo Wagner (27°42'00"S 49°20'02"W), State of Santa Catarina, and the National Park of Aparados da Serra in the municipality of Cambará do Sul (29°08'02"S 49°59'40"W), State of Rio Grande do Sul. Both sites are covered by *Araucaria angustifolia* forests and are located on the South Brazilian Plateau at altitudes of 900–1200 m (Veloso et al. 1991). Köppen (1948) classified the region's climate as Cfb: mesothermic, wet (with regular rainfall throughout the year), and with mild summers.

The specimens were examined morphologically according to Miller & Miller (1988). Colors were coded according to the color chart of Kornerup & Wanscher (1978). Measurements and photographs were taken using a Motic BA310 microscope equipped with a Moticam 2500 digital camera and Motic Image Plus 2.0 software. Scanning electron microscopy (SEM) was performed at the Center of Electronic Microscopy of the Universidade Federal do Rio Grande do Sul (CME/UFRGS) under a Jeol JSM-6060 SEM. All studied specimens are preserved at the Herbarium, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil (ICN). Five collections from Rio Grande do Sul conserved in the ICN Herbarium representing species closest to *M. austromontana* were also examined.

Taxonomy

Morganella austromontana C.R. Alves, Cortez & R.M. Silveira, **sp. nov.** FIGS 1–6


MYCOBANK MB 820972

Differs from *Morganella subincarnata* by its lighter exoperidium, its ovoid basidiospores, and its habitat on soil in araucariaceous forest.

TYPE—Brazil, Santa Catarina State: Alfredo Wagner, Reserva do Rio das Furnas, 7.I.2014, leg. C.R. Alves 94 (Holotype, ICN 181431).

ETYMOLOGY—The name, meaning “from the southern mountains,” refers to the montane type region on the South Brazilian Plateau.

BASIDIOMES 10 mm high, 8–20 mm diam., subglobose to pyriform. EXOPERIDIUM spiny, formed of short spines (<1 mm high), numerous and densely grouped on the top of the basidioma, light brown (6D5) when fresh, falling away as plates detach from the endoperidium when mature. ENDOPERIDIUM papery, smooth to areolate, white (1A1) when fresh, then yellow grey (2B2) after dried. STERILE BASE poorly developed to present, reaching 3 mm high, yellowish grey (2B2), with scattered, white and thin rhizomorphs (<1 mm thickness).



FIGS 1–3. *Morganella austromontana* (holotype ICN 181431). 1. Basidiome in situ. 2. Dried basidiomata. 3. Paracapillitium and basidiospores. Scale bars: 1, 2 = 10 mm; 3 = 10 μ m.

GLEBA cottony, brownish grey (2D2). SUBGLEBA present, ≤ 3 mm high, cellular, greyish yellow (1B3). BASIDIOSPORES $4\text{--}5 \times 3\text{--}4$ μ m, ovoid to subglobose, pale green in 5% KOH, surface verrucose under light microscope, spiny under SEM;

spines cone-shaped, $<2\ \mu\text{m}$ high. PARACAPILLITIUM $7\text{--}10\ \mu\text{m}$ diam., inflated, with numerous small pores c. $1\ \mu\text{m}$ diam., amorphous matter surrounding the hyphae, septa regular, sometimes, with dichotomous tips. EXOPERIDIUM hyphae $10\text{--}26 \times 8\text{--}15\ \mu\text{m}$ diam., globose, subglobose or pyriform, hyaline to pale yellow, walls thin, about $1\ \mu\text{m}$. MYCOSCLEREIDS $25\text{--}80 \times 7\text{--}13\ \mu\text{m}$ diam., irregular in shape, hyaline, walls thick ($3\ \mu\text{m}$) and smooth.


ECOLOGY & DISTRIBUTION—Basidiomata solitary to gregarious, growing on soil, among a few herbs on the trail board in the Mixed Ombrophilous Forest, with *Araucaria angustifolia* (Bertol.) Kuntze (*Araucariaceae*; Paraná pine, Brazilian pine), about $480\text{--}900\ \text{m}$ altitude, during rainy summer. Only known from the South Brazilian Plateau (States of Rio Grande do Sul and Santa Catarina).

ADDITIONAL SPECIMENS EXAMINED — BRAZIL. RIO GRANDE DO SUL STATE: Cambará do Sul, Parque Nacional dos Aparados da Serra, 13.III.2014, leg. A. Gerlach 1486 (ICN 181432); São Francisco de Paula, Veraneio Hampel, 8.I.2006, leg. V.G. Cortez 013/06 (ICN 154553); FLONA, 21.IV.2007, leg. V.G. Cortez 048/07 (ICN 154554); Itaara, Parque Pinhal, 4.V.2007, leg. V.G. Cortez 075/07 (ICN 154555); Santa Maria, FEPAGRO, 15.V.2007, leg. V.G. Cortez s.n (ICN 154556); Campus UFSM, 4.VI.2007, leg. G. Coelho 485-7 (ICN 154557).

COMMENTS—*Morganella austromontana* is described as a new based on its combination of mainly ovoid ornamented basidiospores (verrucose under LM; shortly spiny under SEM) and an exoperidium densely covered with short white spines in younger basidiomata that become pale brownish and fall as plates in maturity. Another diagnostic feature is the relatively wide ($\leq 10\ \mu\text{m}$) diameter of paracapillitium hyphae, which is rarely greater than $5\text{--}6\ \mu\text{m}$ in other species. The thick-walled mycosclereids found in the exoperidium of *M. austromontana* have also been observed in *M. sulcatostoma* C.R. Alves & Cortez (Alves & Cortez 2013).

Morganella austromontana can be nested among members of *Morganella* sect. *Subincarnata* sensu Ponce de Leon (1971), a section that accommodates species with a deciduous exoperidium and reticulate-areolate endoperidium. Within *M. sect. Subincarnata*, *M. costaricensis* M.I. Morales, which shares with *M. austromontana* an exoperidium with deciduous spines and areolate endoperidium, can be separated by its smooth globose to subglobose basidiospores (Morales et al. 1974, Suárez & Wright 1996).

The type species of the section, *Morganella subincarnata* (Peck) Kreisel & Dring, appears related to *M. austromontana* based on its similar pattern of exo- and endoperidium but differs in its darker (olivaceous brown) exoperidium, its globose basidiospores ($3.5 \times 6.5\ \mu\text{m}$ diam.), and its preferred substrate on



FIGS 4–6. *Morganella austromontana* (holotype ICN 181431). 4. Basidiospores and paracapillitium. 5, 6. Basidiospores (SEM). Scale bars 4 = 10 μm ; 5, 6 = 1 μm .

angiospermous wood (Bowerman 1961, as *Lycoperdon subincarnatum*; Coker & Couch 1928, as *Lycoperdon subincarnatum*; Kreisel & Dring 1967; Ponce de Leon 1971).

Another noteworthy feature of *Morganella austromontana* is the fact its basidiomata were collected from soil, instead of rotting wood, the usual substrate for most *Morganella* species. However, a soil substrate is not exclusive to *M. austromontana*; for example, *M. arenicola* Alfredo & Baseia grows on sandy soil (Alfredo et al. 2014).

Further fieldwork will be necessary to determine if the proposed species is, in fact, restricted to the southern Brazilian highlands and, in particular, to *A. angustifolia* forests. However, based on the current available data and ongoing survey of the gasteroid fungi of Southern Brazil, *M. austromontana* appears strongly associated with these environments.

Acknowledgments

We thank the financial support of Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), the reviewers Dr. Maria Luciana H. Caffot and Dr. Iuri G. Baseia for the considerations, Instituto Chico Mendes (ICMBio) for allowing fieldwork, and Alice C. L. Gerlach for specimen collection.

Literature cited

- Alfredo DS, Baseia IG. 2014. *Morganella nuda*, a new puffball (Agaricaceae, Basidiomycota) in the upland forests of the Brazilian semi-arid region. *Nova Hedwigia* 98: 459–466. <http://dx.doi.org/10.1127/0029-5035/2014/0164>
- Alfredo DS, Leite AG, Braga-Neto R, Baseia IG. 2012. Two new *Morganella* species from the Brazilian Amazon rainforest. *Mycosphere* 3: 66–71. <http://dx.doi.org/10.5943/mycosphere/3/1/8>
- Alfredo DS, Accioly T, Baseia IG. 2014. *Morganella arenicola*, a new species record from North and Northeast Brazil. *Turk. J. Bot.* 38: 595–599. <http://dx.doi.org/10.3906/bot-1307-68>
- Alves CR, Cortez VG. 2013. *Morganella sulcatostoma* sp. nov. (Agaricales, Basidiomycota) from Paraná State, Brazil. *Nova Hedwigia* 96: 409–417. <http://dx.doi.org/10.1127/0029-5035/2013/0078>
- Barbosa MMB, Silva MA, Cruz RHSE, Calonge FD, Baseia IG. 2011. First report of *Morganella compacta* (Agaricales, Lycoperdaceae) from South America. *Mycotaxon* 116: 381–386. <http://dx.doi.org/10.5248/116.381>
- Bowerman CA. 1961. *Lycoperdon* in eastern Canada with special reference to the Ottawa district. *Can J Bot* 39: 353–383. <http://dx.doi.org/10.1139/b61-029>
- Coker WC, Couch JN. 1928. The gasteromycetes of the eastern United States and Canada. University of North Carolina Press, Chapel Hill.
- Cortez VG, Calonge FD, Baseia IG. 2007. Rick's species revision 2: *Lycoperdon benjaminii* recombined in *Morganella*. *Mycotaxon* 102: 425–429.
- Köppen WP. 1948. Climatología, con un estudio de los climas de la Tierra. Fondo de Cultura Económica, México.

- Kornerup A, Wanscher JH. 1978. Methuen handbook of colour. 3rd ed. Eyre Methuen, London.
- Kreisel H, Dring DM. 1967. An emendation of the genus *Morganella* Zeller (*Lycoperdaceae*). Feddes Repert. 74: 109–122. <http://dx.doi.org/10.1002/fedr.19670740105>
- Miller OK, Miller HH. 1988. Gasteromycetes: morphological and development features with keys to the orders, families and genera. Mad River, Eureka.
- Morales IM, Nassar M, Saénz JA. 1974. *Lycoperdaceae* of Costa Rica I. The genus *Morganella*. Rev. Biol. Trop. 21: 221–227.
- Ponce de Leon P. 1971. Revision of the genus *Morganella* (*Lycoperdaceae*). Fieldiana, Bot. 34: 27–44. <http://dx.doi.org/10.5962/bhl.title.2567>
- Suárez VL, Wright JE. 1996. South American gasteromycetes V: the genus *Morganella*. Mycologia 88: 655–661. <http://dx.doi.org/10.2307/3761163>
- Trierveiler-Pereira L, Baseia IG. 2009. A checklist of the Brazilian gasteroid fungi (*Basidiomycota*). <http://www.mycotaxon.com/resources/checklists/trierweiler-v108-checklist.pdf> [Summary: Mycotaxon 108: 441–444. <http://dx.doi.org/10.5248/108.441>].
- Veloso HP, Rangel-Filho ALR, Lima JCA. 1991. Classificação da vegetação brasileira adaptada a um sistema universal. IBGE, Rio de Janeiro.
- Zeller SM. 1948. Notes on certain gasteromycetes including two new orders. Mycologia 40: 639–668. <http://dx.doi.org/10.2307/3755316>